

Article

[Prev](#) | [Next](#) | [Table of Contents](#)

Characterization of the Most Odor-Active Compounds in an American Bourbon Whisky by Application of the Aroma Extract Dilution Analysis

Luigi Poisson and Peter Schieberle*

Deutsche Forschungsanstalt für Lebensmittelchemie,
Lichtenbergstrasse 4, D-85748 Garching, Germany

J. Agric. Food Chem., 2008, 56 (14), pp 5813-5819

DOI: 10.1021/jf800382m

Publication Date (Web): June 21, 2008

Copyright © 2008 American Chemical Society

* Corresponding author. Tel: +49 89 289 13265. Fax: +49 89 289 14183. E-mail: Peter.Schieberle@ch.tum.de.

Abstract

Figures

Full Text HTML

References

Hi-Res PDF [205 KB]

PDF w/ Links [173 KB]

Abstract

Application of the aroma extract dilution analysis (AEDA) on the volatile fraction carefully isolated from an American Bourbon whisky revealed 45 odor-active areas in the flavor dilution (FD) factor range of 32–4096 among which (*E*)- β -damascenone and δ -nonalactone showed the highest FD factors of 4096 and 2048, respectively. With FD factors of 1024, (3*S*,4*S*)-*cis*-whiskylactone, γ -decalactone, 4-allyl-2-methoxyphenol (eugenol), and 4-hydroxy-3-methoxy-benzaldehyde (vanillin) additionally contributed to the overall vanilla-like, fruity, and smoky aroma note of the spirit. Application of GC-Olfactometry on the headspace above the whisky revealed 23 aroma-active odorants among which 3-methylbutanal, ethanol, and 2-methylbutanal were identified as additional important aroma compounds. Compared to published data on volatile constituents in whisky, besides ranking the whisky odorants on the basis of their odor potency, 13 aroma compounds were newly identified in this study: ethyl (*S*)-2-methylbutanoate, (*E*)-2-heptenal, (*E,E*)-2,4-nonadienal, (*E*)-2-decenal, (*E,E*)-2,4-decadienal, 2-isopropyl-3-methoxypyrazine, ethyl phenylacetate, 4-methyl acetophenone, α -damascone, 2-phenylethyl propanoate, 3-hydroxy-4,5-dimethyl-2(5*H*)-furanone, trans-ethyl cinnamate, and (*Z*)-6-dodeceno- γ -lactone.

Introduction

Jump to a section

For whisky production, briefly, malt and/or ground cereals are mixed with water to yield a mash that is subsequently fermented with yeast. The resulting beer is then distilled to yield an alcoholic distillate, which is finally stored in barrels (1). It is generally accepted that barrel aging is among the key processing steps in the generation of the unique aroma of whisky.

Typical for American Bourbon whisky is (i) the high content of corn among the cereals used (>51%), (ii) the high alcohol content of the distillate (max. 80% by volume), and (iii) the storage of the distillate in new, heat-charred oak casks. While in the manufacturing of Scottish whisky, the filtered mash (wort) is processed, and the complete mash is fermented in the production of American whiskies. The mash is either fermented with yeast alone (sweet mash) or with a mixture of lactic acid bacteria, and yeast (sour mash) (2). A product stored for at least two years is called straight Bourbon whisky, but Bourbon whisky may not only originate from the Bourbon district in Kentucky but also originate from other areas, if it is manufactured according to the requirements given above (2).

For more than 40 years, investigations on the volatile components of whisky have been performed, and today, more than 300 compounds have been identified in several types of

Related Content

Characterization of the Key Aroma Compounds in an American Bourbon Whisky by Quantitative Measurements, Aroma Recombination, and Omission Studies
Journal of Agricultural and Food Chemistry

Formation of α -Dicarbonyl Compounds in Beer during Storage of Pilsner
Journal of Agricultural and Food Chemistry

Characterization of Odor-Active Compounds in Extracts Obtained by Simultaneous Extraction/Distillation from Moroccan Black Olives
Journal of Agricultural and Food Chemistry

Other ACS articles by these authors:

Luigi Poisson
Peter Schieberle

Article Tools

Add to Favorites

Download Citation

Email a Colleague

Permalink

Rights & Permissions

Citation Alerts

SciFinder Links



View Reference Detail

History

Published In Issue
July 23, 2008Article ASAP
June 21, 2008Received: February 7, 2008
Accepted: April 27, 2008
Revised: April 23, 2008

Recommend & Share

CiteULike

Delicious

Digg This

Facebook

Newsvine

whisky (3-5). Already in 1963, Nykaenen and Suomalainen (6) identified several major components of the volatile fraction of American Bourbon as well as Scottish whisky, such as 2-methylpropanol, 3-methylpropanol, 2-phenylethanol, and acetaldehyde as well as acids and esters.

Steinke and Paulson (7) characterized various phenols in thermally processed grains, and they could show that 4-vinylphenol and 4-vinyl-2-methoxyphenol are formed during distillation by a decarboxylation of the precursors *p*-coumaric acid or ferulic acid, respectively, present in the cereals. The norisoprenoids α - and β -ionone were first detected in Bourbon whisky by LaRoe et al. (8), and α - and β -carotene were suggested as their precursors from corn. Another norisoprenoide, (*E*)- β -damascenone, was later identified in Bourbon whisky by Masuda and Nishimura (9), while Suomalainen and Nykaenen (10) identified the whisky lactone with a coconut-like smell for the first time.

In order to evaluate the contribution of individual volatile components to the aroma of whisky, Salo and co-workers (11, 12) were the first to calculate odor units for individual components on the basis of the ratio of their concentration in whisky and their odor thresholds in an ethanol/water mixture. The authors proposed in particular methylpropanal, butanal, 3-methylbutanal, pentanal, 2,3-butandione, and esters, such as ethyl acetate, ethyl hexanoate, and ethyl octanoate, as important aroma components. In addition, results obtained by preparing a model aroma mixture (12, 13) have led to the assumption that carbonyl compounds and esters are the most important odorants in whisky. Spiking of whisky with the respective reference compounds also revealed the importance of phenol, methylphenols, and 2-methoxyphenol as aroma-active constituents (14, 15).

On the basis of approaches using molecular sensory science, it has been shown for a considerable number of foods that the entire set of volatiles present in a food is not able to interact with human olfactory receptors (16). Instead, only a smaller number of the so-called key odorants is obviously detected by the human odorant receptors and is able to initiate a cascade of biochemical reactions finally leading to aroma perception in the brain.

An approach to separate odor-active volatiles from the bulk of odorless food volatiles is GC-Olfactometry (GC-O) or, more comprehensively, dilution to odor threshold techniques, such as aroma extract dilution analysis (16). However, only a very few studies using such techniques to unravel the key aroma compounds of whisky have been reported to date.

Only Connor et al. (17) applied GC-O on an extract obtained by SPME from Scotch whisky and particularly confirmed the importance of *cis*-whisky lactone and vanillin in the overall aroma of this type of whisky.

In order to systematically improve and/or modify the aroma of whisky by changing the recipe or the manufacturing technology, there is a need to reveal the influence of each technological step on aroma compound formation during whisky processing. But, although numerous investigations have been undertaken to follow changes in selected volatiles caused by processing, in particular, a comprehensive approach aimed at identifying the entire set of odor-active compounds in Bourbon whisky is still lacking. Because the knowledge on the key odorants in the final product is the prerequisite for studies on the influence of processing steps, the aim of the present study was to locate the potent odorants in an aroma extract from a commercial American Bourbon whisky by application of the aroma extract dilution analysis and to identify the most odor-active compounds by means of reference odorants.

Materials and methods

Jump to a section

Whisky

The whisky under investigation was a Kentucky Straight Bourbon whisky which, according to the label, had been produced according to the sour mesh method and had been stored in new, heat-charred oak casks for at least three years. Several batches of the same product (same year of production) were purchased at a local supermarket. Mentioning of a brand name does not imply any research contact with the whisky manufacturer nor is it done for advertising purposes.

Chemicals

Reference compounds of the odorants identified were obtained from the commercial sources given in parentheses: 4-allyl-2-methoxyphenol, *a*-damascone, (*E,E*)-2,4-decadienal, γ -decalactone, 1,1-diethoxyethane, γ -dodecalactone, ethyl acetate, ethyl hexanoate, 4-ethyl-2-methoxyphenol, ethyl (*S*)-2-methylbutanoate, ethyl 3-methylbutanoate, ethyl octanoate, 4-ethylphenol, ethyl 2-phenylacetate, 3-hydroxy-4,5-dimethyl-2(*5H*)-furanone, 2-isopropyl-3-methoxypyrazine, 4-methyl acetophenone, 2-methylbutanal, 3-methylbutanal, 3-methylbutanol, 3-methylbutyl acetate, 2-methylpropanal, (*E,E*)-2,4-nonadienal, (*E,Z*)-2,6-nonadienal, γ -nonalactone, δ -nonalactone, (*E*)-2-nonenal, phenylacetaldehyde, 2-phenylacetic acid, 2-phenylethyl acetate, 2-phenylethyl propanoate, *cis/trans*-whiskylactone, (*E*)-2-decenal, dimethyl sulfide, *trans*-ethyl cinnamate, ethyl butanoate, ethyl 2-methylpropanoate, ethyl pentanoate, ethyl propanoate, (*E*)-2-heptenal, 2-methylpropanol, nonanal, and 2-phenylethanol (Sigma-Aldrich Chemie, Taufkirchen, Germany); acetaldehyde, ethanol, 4-hydroxy-3-methoxybenzaldehyde, and 2-methoxyphenol (Merck, Darmstadt, Germany); β -ionone (Roth, Karlsruhe, Germany); (*E*)- β -damascenone was a gift from Symrise, Holzminden, Germany.

The following odorants were synthesized as reported in the literature given in parentheses: (*Z*)-6-dodeceno- γ -lactone (18) and (*Z*)-2-nonenal (19).

Diethyl ether and *n*-pentane (Merck, Darmstadt, Germany) were freshly distilled before use. Silica 60 (Merck, Darmstadt, Germany) was purified according to a procedure described in ref 20.

Isolation of Volatiles

The volatiles present in whisky (25 mL) were isolated by extraction of a 1:1 dilution with tap water (saturated with NaCl) using diethyl ether (3 \times 100 mL). The combined extracts were dried over anhydrous sodium sulfate and concentrated to 100 mL at 37 °C using a Vigreux column (50 cm \times 1 cm i.d.). The nonvolatile compounds were removed by high vacuum distillation using the SAFE apparatus (21), and the distillate was concentrated to 100 μ L by means of a Vigreux column (60 cm \times 1 cm i.d.) followed by microdistillation (16). The distillate was separated into a fraction containing the neutral/basic volatiles (NBF) and the acidic volatiles (AF) by treatment with an aqueous sodium bicarbonate solution (22).

Fractionation of the Volatiles by Column Chromatography

The NBF fraction obtained from whisky (200 mL) was concentrated to 1 mL and was applied onto a water-cooled glass column (30 cm \times 1 cm) filled with silica 60 (30 g) in *n*-pentane. To yield five fractions containing compounds of increasing polarity, the distillate was separated using five *n*-pentane/diethyl ether mixtures (100 mL each; fraction A, 100:0; fraction B, 95:5; fraction C, 90:10; fraction D, 80:20; fraction E, 50:50 by vol.). Each fraction was dried over anhydrous sodium sulfate and concentrated to 1 mL as described above.

High Resolution Gas Chromatography–Olfactometry (HRGC-O) and Mass Spectrometry (MS)

HRGC was performed by means of a gas chromatograph type 8160 (Fisons Instruments, Mainz, Germany) with helium as the carrier gas at a pressure of 70 kPa. Samples were applied by the cold-on-column-injection technique onto either capillary DB-5 (CP-Sil 8CB) (Chrompack, Frankfurt, Germany), DB-FFAP, or DB-1701 (J & W Scientific; supplied by Agilent Technologies) (Waldbronn, Germany). The dimensions for all three columns were 30 m length, 0.32 mm i.d., and 0.25 μ m film thickness. For GC-Olfactometry, the end of the capillary was connected to a deactivated Y-shaped glass splitter dividing the effluent of the column into two equal parts, which was then transferred via two deactivated but uncoated fused silica capillaries (50 cm \times 0.25 mm) to a sniffing port and an FID, respectively. The sniffing port, which was mounted on a detector base of the GC, was held at 180 °C, and the FID was operated at 240 °C. On-column injection of the samples (0.5 μ L) was performed at an oven temperature of 35 °C. After 2 min, the temperature was raised by 40 °C per min to 50 °C, held for 2 min, and raised by 6 °C per min to 180 °C. Then, the temperature was raised by 20 °C per min to 240 °C and finally held for 10 min. During a GC run, a panelist placed his/her nose close to and above the top of the sniffing port and evaluated the odor of the chromatographic effluent. All compounds detected during the evaluation of the extract by four trained panelists were considered relevant, and no anosmic panelist was used.

Linear retention indices (RI) of the compounds were calculated from the retention times of *n*-

alkanes (C-8 to C-22) (22). Mass spectra were recorded using a gas chromatograph 5890 series II (Hewlett-Packard, Waldbronn, Germany) connected to a sector field mass spectrometer type MAT 95 S (Finnigan, Bremen, Germany). Mass spectra in the electron ionization mode (MS-EI) were recorded at 70 eV ionization energy and mass spectra in the chemical ionization mode (MS-CI) at 115 eV using isobutane as the reactant gas.

Aroma Extract Dilution Analysis (AEDA)

For AEDA, fractions containing the neutral/basic (NBF) and acidic volatiles (AF) were stepwise diluted using diethyl ether as the solvent to obtain dilutions of 1:1, 1:2, 1:4, 1:8, 1:16... 1:4096 of the original extracts (16). Sniffing of both series of dilutions was continued until no odorant could be detected by GC-O. Each odorant was thus assigned a flavor dilution factor (FD factor) representing the last dilution in which the odorant was still detectable. The evaluation of the original extract was performed by four panelists, in particular, to agree upon the odor-active areas detectable and also upon their odor qualities. Odor qualities were assigned on the basis of a flavor language developed in our group by an evaluation of the odor quality of reference odorants (Czerny et al., unpublished work). The AEDA of both fractions was finally performed by two panelists in separate runs, and the FD factors determined were averaged. Training of the panelists was done by GC-O using at least 40 odor-active reference compounds in a concentration 5 times above their odor thresholds in air.

Static Headspace-Olfactometry (SHO) and Aroma Dilution Analysis (ADA)

Aliquots of the whisky (5 mL) were pipetted into septum-sealed vessels (200 mL) and equilibrated for 30 min at room temperature. Decreasing volumes (10 to 0.25 mL) of headspace samples were taken from separate vessels using a gastight syringe (Hamilton, Australia) and were analyzed by HRGC-O as described previously (22).

Determination of Enantiomeric Ratios

The enantiomeric ratios were determined by two-dimensional gas chromatography–mass spectrometry using a chiral column (BGB-176, 30 m × 0.25 mm, film thickness 0.25 μm; BGB Analytik AG, Rothenfluh, Switzerland) in the second dimension. The intensities of the molecular ions (MS-CI) were recorded for the calculation of the enantiomeric ratio (22).

Results and Discussion

Jump to a section

Identification of Odor-Active Compounds

The Bourbon whisky under investigation exhibited an overall vanilla-like fruity aroma with additional smoky, malty odor qualities. By solvent extraction and SAFE distillation (21), a volatile fraction was obtained, which elicited the typical aroma of the whisky when sniffed by a group of 10 panelists on a strip of filter paper in an ethanol/water mixture (4:6 by vol.) equal to the original whisky. Thus, the method of volatile isolation was proven to be appropriate for aroma isolation from whisky.

To obtain insight into the odorants contributing to the overall aroma, an aliquot of the distillate representing the total volatiles of 200 μL of whisky were separated by GC, and the effluent was evaluated by GC-Olfactometry. A total of 45 odor-active areas were detected, and a quite intense cooked-apple-like odor was smelled at position 24 (Figure 1). In addition, several coconut-like odors, 27, 31, and 42, appeared with high intensities. By application of the aroma extract dilution analysis (AEDA), the cooked-apple-like odorant 24, followed by the coconut smelling compound, 31 appeared with the highest flavor dilution (FD) factors (Figure 1). In addition, the vanilla-like smelling odorant 45, a flowery smelling odorant 43, the clove-like smelling odorant 36, the peach-like smelling odorant 35, and the coconut-like smelling odorant 30 showed somewhat lower FD factors.




Figure 1. Gas chromatogram (A) and flavor dilution (FD) chromatogram (B) of the volatile fraction isolated from Bourbon whisky.

To identify the compounds responsible for these odors, first, the retention indices of the odor-

active areas and the respective aroma attributes were compared to data available in an in-house database containing about 500 compounds previously identified by our group as key food odorants. This procedure suggested a structure for most of the aroma compounds. To confirm these structures, a distillate obtained from 200 mL of whisky was separated into five fractions by column chromatography on silica gel, the odor-active areas were relocated in the fractions by GC-O and, then mass spectra in the electron impact mode (MS-EI) and in the chemical ionization mode (MS-CI) were recorded. The data were finally compared to data obtained by the analysis of the respective reference compounds.

Following this procedure, compound **24** with the highest FD factor was identified as (*E*)- β -damascenone (**Figure 2**) exhibiting an intense aroma like that of cooked apples. With compounds **21** and **29** (FD factors of 512 and 128, respectively), two further norisoprenoids were identified, namely, α -damascone, likewise exhibiting a cooked apple-like smell, and β -ionone exhibiting a violet-like odor (**Table 1**). Compound **31** with the second highest FD factor of 2048 was identified as γ -nonalactone (**Figure 2**). The two possible enantiomers were separated on a chiral cyclodextrin stationary phase, and the enantiomeric distribution was determined to be 56% *R*- γ -nonalactone and 44% *S*- γ -nonalactone (data not shown). Two further compounds with a coconut-like odor could be identified as *cis*- and *trans*-whiskylactones (β -methyl- γ -octalactone; **27** and **30**, **Table 1**). Whiskylactone (**30**; **Figure 1**) has two chiral centers at positions 3 and 4 resulting in 4 stereoisomers. Mosandl and co-workers (**23, 24**) have previously shown that only the (3*S*,4*S*)-*cis*-whiskylactone and the (3*S*,4*R*)-*trans*-whiskylactone occur in whisky. Thus, by comparison of the retention indices with published data (**23**), compound **27** was identified as (3*S*,4*R*)-*trans*-whiskylactone and compound **30** as (3*S*,4*S*)-*cis*-whiskylactone. With their high FD factors of 1024 and 512, respectively (**Table 1**), both enantiomers are likely to contribute to the overall whisky aroma.




Figure 2. Structures of the most odor-active compounds (FD \geq 1024) identified in Bourbon whisky (numbering refers to **Table 1**).

Table 1. Most Odor-Active (FD \geq 32) Volatile Constituents Identified in Bourbon Whisky

no.	odorant	odor quality	fract.	RI on				FD factor	earlier reported as volatile compound in whisky
				FFAP	DB-5	1701	DB-1701		
1	1,1-diethoxyethane	fruity	A	900	730	<800	128	(6, 25)	
2	ethyl 2-methylpropanoate	fruity	A	958	756	811	64	(10, 26)	
3	ethyl butanoate	fruity	A	1029	805	857	64	(10, 27)	
4	ethyl (<i>S</i>)-2-methylbutanoate	fruity	A	1040	850	900	256		
5	ethyl 3-methylbutanoate	fruity	A	1049	849	906	32	(3)	
6	2-methylpropan-1-ol	malty	C	1091	648	n.d.	32	(6, 25)	
7	unknown	fruity	B	1102	988	n.d.	32		
8	3-methylbutyl acetate	fruity	B	1120	879	948	256	(6, 25)	
9	3-methylbutan-1-ol	malty	C	1215	735	844	512	(6, 25)	
10	ethyl hexanoate	fruity	B	1218	1000	1062	128	(6, 10)	
11	(<i>E</i>)-2-heptenal	fatty, green	B	1311	959	1063	256		
12	nonanal	soapy	B	1380	1100	n.d.	32	(28)	
13	2-isopropyl-3-methoxypyrazine	earthy	C	1413	1091	1145	32		
14	ethyl octanoate	fruity	B + C	1420	1197	1263	128	(6, 25)	
15	(<i>E</i>)-2-nonenal	green	C + D	1533	1159	1275	128	(29)	
16	(<i>E,Z</i>)-2,6-nonadienal	green	C + D	1578	1153	n.d.	256	(30)	
17	(<i>E</i>)-2-decenal	fatty	D	1624	1260	1370	32		

17	(E)-2-decenal	fatty	D	1024	1260	1370	32	
18	(E,E)-2,4-nonadienal	fatty	D	1691	1211	n.d.	32	
19	ethyl 2-phenylacetate	flowery	D	1695	n.d.	n.d.	128	
20	4-methylacetophenone	sweet, almond-like		1747	1186	1326	64	
21	α -damascone	cooked apple	C	1779	1389	n.d.	512	
22	(E,E)-2,4-decadienal	fatty	B + C	1791	1315	1447	128	
23	2-phenylethyl acetate	flowery	B	1804	1256	1371	128	(10, 25)
24	(E)- β -damascenone	cooked apple	B	1813	1389	1487	4096	(9)
25	2-methoxyphenol	phenolic	C	1849	1089	1225	64	(3, 31)
26	2-phenylethyl propanoate	fruity		1860	n.d.	n.d.	128	
27	(3S,4R)- <i>trans</i> -whiskylactone	coconut-like	C	1866	1295	1487	512	(31, 32)
28	2-phenylethanol	flowery	D	1900	1113	1273	512	(6, 25)
29	β -ionone	violet-like	C	1929	1495	1620	128	(8)
30	(3S,4S)- <i>cis</i> -whiskylactone	coconut-like	C + D	1946	1328	1522	1024	(31, 32)
31	(R/S)- γ -nonalactone	coconut-like	D	2018	1365	1566	2048	(26, 31)
32	4-ethyl-2-methoxyphenol	phenolic, clove-like	D	2030	1284	1413	128	(7, 31)
33	δ -nonalactone	peach-like	D	2047	n.d.	1612	256	(15, 26)
34	<i>trans</i> ethyl cinnamate	fruity	D	2108	1461	n.d.	256	
35	γ -decalactone	peach-like	D	2132	1471	1690	1024	(5)
36	4-allyl-2-methoxyphenol	clove-like	C	2161	1357	1498	1024	(31)
37	4-ethylphenol	phenolic	C	2168	1168	1395	32	(7, 31)
38	3-hydroxy-4,5-dimethyl-2(5H)-furanone	seasoning-like	D	2206	1100	1362	128	
39	unknown	coconut-like	D	2243	1562	1805	512	
40	unknown	fruity	D	2265	n.d.	n.d.	128	
41	γ -dodecalactone	peach-like	D	2381	1685	1900	256	(5)
42	(Z)-6-dodeceno- γ -lactone	peach-like	D	2425	1680	n.d.	512	
43	unknown	flowery	A + E	2442	n.d.	n.d.	1024	
44	2-phenylacetic acid	flowery	E	2552	1275	n.d.	256	(33)
45	4-hydroxy-3-methoxy-benzaldehyde	vanilla-like	D	2600	1400	1632	1024	(34)

a Numbering refers to **Figure 1**.

b The odorant was identified by comparing it with the reference compound on the basis of the following criteria: (i) retention indices on the capillaries detailed in the table, (ii) mass spectra obtained by MS-EI and MS-CI, and (iii) odor quality and odor threshold determined by GC-Olfactometry.

c Odor quality perceived at the sniffing port.

d The odorant was detected in the column chromatography fractions A-E.

e Retention index; n.d. = not determined.

f Because no unequivocal mass spectrum could be obtained, the odorant was identified by the remaining criteria given in footnote b.

Compound **36** with a clove-like, phenolic odor and a high FD factor of 1024 was identified as 4-allyl-2-methoxyphenol (**Figure 2**). Further odorants showing phenolic odor attributes, but with somewhat lower FD factors, were identified as 4-ethyl-2-methoxyphenol (**32**; **Table 1**) and 2-methoxyphenol (**25**; **Table 1**). Among these compounds having the same ortho-methoxy-hydroxy substitution at the aromatic ring, which were probably extracted from the degraded lignin of the oak cask, 4-hydroxy-3-methoxybenzaldehyde (Vanillin; **45**; **Figure 2**) with a vanilla-like aroma showed the highest FD factor.

The largest peak in the gas chromatogram (9; Figure 1) showing a malty smell was identified as 3-methylbutanol, and the aroma-active area 28 exhibiting a flowery aroma note was identified as 2-phenylethanol. For both alcohols, well known as metabolites of amino acid degradation by yeasts, an FD factor of 512 was determined.

Various esters exhibiting fruity or flowery aroma notes, were identified in the whisky distillate (Table 1), namely, ethyl 2-methylbutanoate (4), 3-methylbutyl acetate (8), ethyl hexanoate (10), ethyl methylpropanoate (2), ethyl butanoate (3), 2-phenylethyl acetate (23), 2-phenylethyl propanoate (26), *trans*-ethyl cinnamate (34), and ethyl 2-phenylacetate (19). Ethyl 2-methylbutanoate (4) was shown to be present as a nearly pure (*S*)-enantiomer (98%) when analyzed on a chiral cyclodextrin phase (data not shown). Ethyl (*S*)-2-methylbutanoate (4), ethyl 2-phenylacetate (19), and *trans*-ethyl cinnamate (34) were all identified for the first time in whisky.

Several aldehydes, exhibiting fatty or green odor notes, were also identified for the first time in the volatile fraction of whisky, among them were (*E*)-2-heptenal and (*E*)-2-decenal (Table 1). The results of the identification experiments in combination with the FD factors are summarized in Table 1, which also shows, whether a compound has previously been reported as constituent of whisky.

The extraction and distillation/concentration process used in the AEDA experiments, may lead to losses in highly volatile compounds. To overcome this gap, static headspace-olfactometry (SHO) was applied to the whisky sample. The results (Table 2) revealed 23 aroma-active areas, and the aroma contribution of these highly volatile components was evaluated by analyzing decreasing volumes of the headspace above the whisky. In the largest volume of 10 mL, 23 odor-active areas were detected. By decreasing the volume down to 100 μ L, only the malty smelling 3-methylbutanal could finally be detected, corresponding to a relative FD factor of 100. Further odorants with a high FD factor were identified as ethanol and ethyl 2-methylpropanoate.

Table 2. Most Odor-Active Volatile Constituents in the Headspace above the Bourbon Whisky

odorant	odorquality	RI onSE-54	vol. [mL]	RDF-factor	earlier reported as volatile compound in whisky
acetaldehyde	solvent-like	<600	0.5	20	(6)
dimethylsulfide	canned maize	<600	10.0	1	(35)
ethanol	ethanolic	<600	0.25	40	(6)
ethyl acetate	fruity	<600	2.5	4	(6)
2-methylpropanal	malty	562	5.0	2	(26)
3-methylbutanal	malty	650	0.1	100	(26)
2-methylbutanal	malty	660	0.5	20	(26)
ethyl propanoate	fruity	711	2.5	4	(27)
1,1-diethoxyethane	fruity	726	0.5	20	(6)
3-methylbutanol	malty	733	0.5	20	(6)
ethyl 2-methyl propanoate	fruity	754	0.25	40	(26)
unknown	fruity	757	0.25	40	
unknown	sulfury	771	0.5	20	
ethyl butanoate	fruity	803	1.0	10	(36)
unknown	earthy	818	1.0	10	
ethyl (<i>S</i>)-2-methylbutanoate	fruity	846	0.5	20	
ethyl 3-methylbutanoate	fruity	848	5.0	2	(3)
3-methylbutyl acetate	fruity	890	2.5	4	(6)
ethyl pentanoate	fruity	904	5.0	2	(26)
unknown	roasty	926	5.0	2	
unknown	green	958	5.0	2	
unknown	fruity	971	2.5	4	
ethyl hexanoate	fruity	1000	2.5	4	(6)

a See Table 1.

b See Table 1.

c Retention index.

d Smallest headspace volume in which the odorant was detectable.

e The relative flavor dilution factor (RDF) was calculated by dividing the largest headspace volume analyzed (10 mL) by the smallest volume in which the odorant was detectable by GC-O.

Although in our study no experiments were performed to identify precursors of the key odorants detected in whisky, several suggestions on their origin are given on the basis of literature data reported for other alcoholic beverages: (*E*)- β -damascenone, which appeared with the highest FD factor has already been detected in various alcoholic beverages, for example, rum, beer, and wine (37-39). In whisky, it has been reported for the first time by Masuda and Nishimura (9). In studies on the formation of (*E*)- β -damascenone, Roberts and Acree (40) suggested the acid-catalyzed formation of (*E*)- β -damascenone from 3-hydroxy-7,8-dihydro- β -ionol as the key precursor. This precursor has also been identified in wine (41, 42). On the basis of these findings, it can be assumed that cereals may contain similar precursors leading to the generation of (*E*)- β -damascenone during fermentation and/or distillation.

Whiskylactone was first identified in whisky by Nishimura and Masuda in 1971 (31) and has been the subject of investigation regarding whisky flavor ever since. Mosandl et al. (23, 24) reported that both enantiomers (3*S*,4*S*)-*cis*- and (3*S*,4*R*)-*trans*-whiskylactone are derived from the wood of American white oak (*Quercus alba*). Previously 3-methyl-4-(3,4-dihydro-5-methoxybenzoyloxy)-octanoic acid has been identified in extracts from *Quercus alba* and has been postulated as the precursor of the lactones during charring of barrels (43, 44). Consequently, Connor et al. (17) found that whisky lactone is present in significantly higher amounts in whisky that was stored in new, heat-treated oak casks as compared to whisky that was aged in re-used casks.

(*E*,*Z*)-2,6-Nonadienal was previously identified in malt whisky by Wanikawa et al. (30). During the process of malting, linoleic and linolenic acid present in cereals are suggested to be oxidized by lipoxygenases into hydroperoxides, which might be decomposed into unsaturated aldehydes such as (*E*)-2-nonenal. Linoleic acid can also be formed during yeast fermentation (45) or can be released from oak wood (46). (*E*,*E*)-2,4-Nonadienal, (*E*)-2-decenal, and (*E*)-2-heptenal, identified here for the first time in whisky, may be generated by similar mechanisms of lipid oxidation.

In the present study, a total of 13 aroma compounds were newly detected in the volatile fraction of Bourbon whisky, namely, (*S*)-ethyl 2-methylbutanoate, (*E*)-2-heptenal, (*E*,*E*)-2,4-nonadienal, (*E*)-2-decenal, (*E*,*E*)-2,4-decadienal, 2-isopropyl-3-methoxypyrazine, ethyl phenylacetate, 4-methylacetophenone, α -damascone, 2-phenylethyl propanoate, 3-hydroxy-4,5-dimethyl-2(⁵H)-furanone, *trans*-ethyl cinnamate, and (*Z*)-6-dodeceno- γ -lactone. Previous studies have emphasized the importance of straight-chain ethyl esters, such as ethyl acetate, ethyl butanoate, ethyl hexanoate, and ethyl octanoate for whisky aromas as these are present in whisky in high concentrations (47). Our investigations have, however, shown that, in particular, branched ethyl esters such as ethyl (*S*)-2-methylbutanoate, ethyl 3-methylbutanoate, and ethyl 2-methylpropanoate contribute to the fruity attribute in whisky flavor among which ethyl (*S*)-2-methylbutanoate is suggested as the most important fruity smelling odorant.

To confirm the aroma contribution of the compounds listed in Table 1, quantitative studies and aroma recombination experiments are underway (48). However, the compounds identified can already be suggested as indicators to assess the odor quality of Bourbon whisky.

Acknowledgment

Jump to a section

We thank Dr. Petra Steinhaus for her valuable support in the preparation of this manuscript.


References

Jump to a section

This article references 48 other publications.

1. Watson, D. C. Factors Influencing the Congener Composition of Malt Whisky New Spirit In: *Flavour of Distilled Beverages: Origin and Development*; Piggott, J. R. Ed.; Ellis Horwood: Chichester, England, **1983**, pp 79–92
2. Postel, W.; Adam, L. Characterization of American and Canadian whiskies by their volatile profile (in German) *Branntweinwirtschaft* **1979** 119 172 176 [[ChemPort](#)]
3. Demyttenaere, J. C. R.; Sanchez Martinez, J. K.; Verhe, R.; Sandra, P.; De Kimpe, N. Analysis of volatiles of malt whisky by solid-phase microextraction and stir bar sorptive extraction *J. Chromatogr., A* **2003** 985 221 232 [[CrossRef](#)], [[PubMed](#)], [[ChemPort](#)]
4. Camara, J. S.; Marques, J. C.; Perestrelo, R. M.; Rodrigues, F.; Oliveira, L.; Andrade, P.; Caldiera, M. Comparative study of the whisky aroma profile based on headspace solid phase microextraction using different fibre coating *J. Chromatogr., A* **2007** 1150 198 207 [[CrossRef](#)], [[PubMed](#)], [[ChemPort](#)]
5. The Netherlands Organisation for Applied Scientific Research TNO: Volatile Compounds in Food, Version 9.2. <http://www.vcf-online.nl> (**2007**).
6. Nykänen, L.; Suomalainen, H. The aroma compounds of alcoholic beverages *Teknillisen Kemian Aikakauslehti* **1963** 20 789 795
7. Steinke, R.; Paulson, M. C. The production of steam-volatile phenols during the cooking and alcoholic fermentation of grain *J. Agric. Food Chem.* **1964** 12 381 387 [[ACS Full Text](#) , [[ChemPort](#)]
8. LaRoe, E. G.; Shipley, P. A. Whiskey composition: formation of alpha- and beta ionone by the thermal decomposition of beta-carotene *J. Agric. Food Chem.* **1970** 18 174 175 [[ACS Full Text](#) , [[ChemPort](#)]
9. Masuda, M.; Nishimura, K. Occurrence and formation of damascenone in alcoholic beverages *J. Food Sci.* **1980** 45 396 397 [[CrossRef](#)], [[ChemPort](#)]
10. Suomalainen, H.; Nykaenen, L. Composition of whisky flavour *Proc. Biochem.* **1970** 5 13 18 [[ChemPort](#)]
11. Salo, P.; Nykaenen, L.; Suomalainen, H. Odor thresholds and relative intensities of volatile aroma components in an artificial beverage imitating whisky *J. Food Sci.* **1972** 37 394 398 [[CrossRef](#)], [[ChemPort](#)]
12. Salo, P. Interactions among volatile aroma compounds in a whiskey imitation *Lebensm. Wiss. Technol.* **1973** 6 52 58 [[ChemPort](#)]
13. Jounela-Eriksson, P. Sensory Aspects of Flavor in Alcoholic Beverages. The Strength and Quality of Whisky Aroma. Proceedings of the VIth International Symposium on Olfaction and Taste; Le Magnen, J.; MacLeod, P., Eds.; Information Retrieval Ltd. , London, **1977**, pp 409–419
14. Swan, J. S.; Burtles, S. M. The development of flavour in potable spirits *Chem. Soc. Rev.* **1978** 7 201 211 [[CrossRef](#)], [[ChemPort](#)]
15. Jounela-Eriksson, P.; Lehtonen, M. Phenols in the Aroma of Distilled Beverages. In The Quality of Foods and Beverages. Proceedings of the 2nd International Flavor Conference, Greece; Charalambous, G., Ed. Inglett: New York, **1981**; pp 167–181
16. Schieberle, P. Recent Developments in Methods for Analysis of Flavor Compounds and Their Precursors In *Characterization of Food: Emerging Methods*; Goankar, A. Ed.; Elsevier: Amsterdam, The Netherlands, **1995**; pp 403–431 [[CrossRef](#)]
17. Connor, J.; Reid, K.; Richardson, G. SPME Analysis of Flavor Components in the Headspace of Scotch Whiskey and Their Subsequent Correlation with Sensory Perception

In *Gas Chromatography-Olfactometry: State of the Art*; Leland, J. V.; Schieberle, P. Eds.; ACS Symposium Series 782; American Chemical Society: Washington DC, 2001; pp 113–121

18. Widder, S.; Sen, A.; Grosch, W. Changes in the flavour of butter oil during storages. Identification of potent odorants *Z. Lebensm.-Unters.-Forsch.* **1991** 193 32 35 [[CrossRef](#)], [[ChemPort](#)]
19. Ullrich, F.; Grosch, W. Flavor deterioration of soybean oil: identification of intense odor compounds formed during flavor reversion *Fett Wiss. Technol.* **1988** 90 332 336 [[CrossRef](#)], [[ChemPort](#)]
20. Esterbauer, H. On the autoxidation of methyl linoleate in water *Fette, Seifen, Anstrichm.* **1968** 70 1 4 [[CrossRef](#)], [[ChemPort](#)]
21. Engel, W.; Bahr, W.; Schieberle, P. Solvent assisted flavour evaporation  a new and versatile technique for the careful and direct isolation of aroma compounds from complex food matrices *Eur. Food Res. Technol.* **1999** 209 237 241 [[CrossRef](#)], [[ChemPort](#)]
22. Hinterholzer, A.; Schieberle, P. Identification of the most odor-active volatiles in fresh, hand-extracted juice of valencia late oranges by odor dilution techniques *Flavour Fragr. J.* **1998** 13 49 55 [[CrossRef](#)], [[ChemPort](#)]
23. Günther, C.; Mosandl, A. Stereoisomeric aroma compounds. XV. Chiroselective analysis of natural odorants: Whisky lactone (in German) *Z. Lebensm.-Unters.-Forsch.* **1987** 185 1 4 [[CrossRef](#)]
24. Mosandl, A.; Kustermann, A. Stereoisomeric flavor compounds. XXX. HPLC analysis of chiral γ -lactones from beverages and fruit preparations *Z. Lebensm.-Unters.-Forsch.* **1989** 189 212 215 [[CrossRef](#)], [[ChemPort](#)]
25. Duncan, R. E. B.; Philip, J. M. Methods for the analysis of Scotch whisky *J. Sci. Food Agric.* **1966** 17 208 214 [[CrossRef](#)], [[ChemPort](#)]
26. Kahn, J. H.; Shipley, P. A.; LaRoe, E. G.; Conner, H. A. Whiskey composition: Identification of additional components by gas chromatography-mass spectrometry *J. Food Sci.* **1969** 34 587 591 [[CrossRef](#)], [[ChemPort](#)]
27. Kahn, J. H.; LaRoe, E. G.; Conner, H. A. Whiskey composition: Identification of components by single-pass chromatography-mass spectrometry *J. Food Sci.* **1968** 33 395 400 [[CrossRef](#)], [[ChemPort](#)]
28. Ter Heide, R. Recent Advances in the Knowledge of Alcoholic Beverages. In Colloque international sur les Aromes Alimentaires, Proceedings of the International Symposium on Food Flavors, Lavoisier, Paris, **1982**; pp 27–45
29. Perry, D. R. Odour Intensities of Whisky Compounds In Distilled Beverage Flavour, Piggott, J. R.; Paterson, A. Eds.; Ellis Horwood, Series in Food Science and Technology; VCH: Weinheim, Germany, **1989**; pp 151–169
30. Wanikawa, A.; Hosoi, K.; Kato, T.; Nakagawa, K. Identification of green note compounds in malt whisky using multidimensional gas chromatography *Flavour Fragr. J.* **2002** 17 207 211 [[CrossRef](#)], [[ChemPort](#)]
31. Nishimura, K.; Masuda, M. Minor constituents of whisky fusel oils. I. Basic, phenolic and lactonic compounds *J. Food Sci.* **1971** 36 819 822 [[CrossRef](#)], [[ChemPort](#)]
32. Otsuka, K.; Zenibayashi, Y.; Itoh, M.; Totsuka, A. Presence and significance of two diastereomers of β -methyl- γ -octalactone in aged distilled liquors *Agric. Biol. Chem.* **1974** 38 485 490 [[ChemPort](#)]
33. MacNamara, K.; Brunerie, P.; Squarcia, F.; Rozenblum, A. Investigation of Flavour Compounds in Whiskey Spent Lees. In *Food Flavors: Generation, Analysis and Process Influence*; Charalambous, G. Eds.; Elsevier Science: New York, **1995**; pp 1753–1766 [[CrossRef](#)]

34. Baldwin, S.; Black, R. A.; Andreassen, A. A.; Adams, S. L. Aromatic congener formation in maturation of alcoholic distillates *J. Agric. Food Chem.* **1967** 15 381 385 [[ACS Full Text](#) , [ChemPort](#)]
35. Masuda, M.; Nishimura, K. Changes in volatile sulfur compounds of whisky during aging *J. Food Sci.* **1981** 47 101 105 [[CrossRef](#)], [[ChemPort](#)]
36. Nykaenen, L. Formation and occurrence of flavour compounds in wine and distilled beverages *Am. J. Enol. Viticult.* **1986** 37 84 96
37. Lehtonen, M.; Suomalainen, H. The analytical profile of some whisky brands *Proc. Biochem.* **1979** 14 5 9 [[ChemPort](#)]
38. Schieberle, P. Primary odorants of pale lager beer *Z. Lebensm.-Unters.-Forsch.* **1991** 193 558 565 [[CrossRef](#)], [[ChemPort](#)]
39. Guth, H. Identification of character impact odorants of different white wine varieties *J. Agric. Food Chem.* **1997** 45 3022 3026 [[ACS Full Text](#) , [ChemPort](#)]
40. Roberts, D. D.; Acree, T. E. Developments in the Isolation and Characterization of β -Damascenone Precursors from Apples. In *Fruit Flavors, Biogenesis, Characterization and Authentication* Rouseff, R. L.; Leahy, M. M. Eds.; ACS Symposium Series 596; American Chemical Society: Washington DC, **1995**; pp 190–199
41. Winterhalter, P.; Sefton, M. A.; Williams, P. J. Two-dimensional GC-DCCC analysis of the glycoconjugates of monoterpenes, norisoprenoids and shikimate-derived metabolites from Riesling wine *J. Agric. Food Chem.* **1990** 38 1041 1048 [[ACS Full Text](#) , [ChemPort](#)]
42. Baderscheider, B.; Skouroumounis, G.; Winterhalter, P. Isolation of two glucosidic precursors of β -damascenone from Riesling wine *Nat. Prod. Letters* **1997** 10 111 114 [[CrossRef](#)]
43. Otsuka, K.; Sato, K.; Yamashita, T. Studies on the mechanism of aging of distilled liquors. VIII. Structure of a precursor of β -methyl- γ -octalactone, an aging flavor compound of distilled liquors *J. Ferment. Technol.* **1980** 58 395 398 [[ChemPort](#)]
44. Masson, G.; Guichard, E.; Fournier, N.; Puech, J-L. Stereoisomers of β -methyl- γ -octalactone. II. Contents in the wood of French (*Quercus petraea*) and American (*Quercus alba*) oaks *Am. J. Enol. Viticult.* **1995** 46 424 428 [[ChemPort](#)]
45. De Rijke, D.; TerHeide, R. Flavour compounds in Rum, Cognac and Whisky In *Flavour of Distilled Beverages: Origin and Development*. Piggott, J. R. Eds.; Ellis Horwood: Chichester, **1983**; pp 192–202
46. Reazin, G. H. Chemical mechanisms of whisky maturation *Am. J. Enol. Viticult.* **1981** 32 283 289 [[ChemPort](#)]
47. Ter Heide, R. The Flavour of Distilled Beverages. In *Developments in Food Science 3B (Food Flavours, Part B, The Flavour of Beverages)*; Morton, I. D., MacLeod, A. J., Eds.; Elsevier: Amsterdam, **1986**; pp 239–336
48. Poisson, L.; Schieberle, P. Characterization of the key aroma compounds in an American bourbon whisky by quantitative measurements, aroma recombination and omission studies. *J. Agric. Food Chem.* **2008**, 5820–5826. [[ACS Full Text](#) , [PubMed](#)]